



# FLUXYR<sup>®</sup> 200 EC

## HERBICIDE

### PCS 05787

An emulsifiable concentrate containing 200g/L fluroxypyr  
(equivalent to 28.6% w/w fluroxypyr-1-methylheptyl ester with an aromatic hydrocarbon)

For the control of certain annual weeds in the listed cereals, maize and grassland.

FOR PROFESSIONAL USE ONLY



### Danger

**Flammable liquid and vapour.**

**May be fatal if swallowed and enters airways.**

**Causes serious eye irritation.**

**May cause respiratory irritation.**

**May cause drowsiness or dizziness.**

**Very toxic to aquatic life with long lasting effects.**

Keep away from heat/sparks/open flames/hot surfaces.

Avoid breathing fume/gas/mist/vapours/spray.

Wear protective gloves/protective clothing/eye protection/face protection.

IF SWALLOWED: Rinse mouth. Do not induce vomiting.

IF INHALED: Remove victim to fresh air and keep at rest in a position comfortable for breathing.

IF IN EYES: Rinse cautiously with water for several minutes.

Remove contact lenses if present and easy to do – continue rinsing.

If eye irritation persists get medical advice/attention.

Collect spillage. Hazardous to the aquatic environment.

Store in a well ventilated place. Keep container tightly closed.

Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for triple-rinsed empty containers which can be disposed of as non-hazardous waste.

**Repeated exposure may cause skin dryness and cracking.**

**To avoid risks to human health and the environment, comply with the instructions for use.**

PCS 05787

## DIRECTIONS FOR USE

| <b>Crop</b>                        | <b>Maximum individual dose (L/product/ha)</b> | <b>Maximum total dose (L/product/ha)</b> | <b>Latest time of application</b>          |
|------------------------------------|---|--|--|
| Spring wheat, spring barley        | 0.75  | 0.75                                     | Flag leaf ligule just visible (GS39)       |
| Winter wheat, winter barley        | 1.0   | 1.0                                      | Flag leaf ligule just visible (GS39)       |
| Winter rye, winter oats, triticale | 1.0   | 1.0                                      | Before second node detectable stage (GS32) |
| Spring oats                        | 0.75  | 0.75                                     | Before second node detectable stage (GS32) |
| Forage maize                       | 1.0   | 1.0                                      | Before 7 leaves unfolded (GS17)            |
| Grassland (established)            | 2.0   | 2.0                                      | 7 days livestock exclusion                 |
| Grassland (new leys)               | 0.75  | 0.75                                     | 7 days livestock exclusion                 |

### **Other specific restrictions**

A maximum total dose of 0.75 litres per hectare must be observed for applications made to cereals between crop emergence in the year of planting and 1st February in the year of harvest.

When the product is applied as a spot treatment to permanent grassland and rotational grass, the maximum concentration must not exceed 30ml product per 10 litres water.

To protect aquatic organisms respect an unsprayed buffer zone of 5m to surface water bodies.

### **CONDITIONS OF APPLICATION**

**CEREALS:** Avoid spraying during periods of prolonged cold (temperatures below 10°C).

**MAIZE:** Avoid spraying during periods where night temperatures are below 5°C.

### **SPECIFIC RECOMMENDATIONS**

**WARNING:** Danger to neighbouring broadleaved crops. Avoid spraying near broadleaved crops in windy conditions.

Use as low a spray pressure as possible and the lowest boom height possible, whilst ensuring an even spray. Do not apply when temperatures exceed 25°C in the shade.

Do not apply FLUXYR 200 EC if the field is undersown with legumes or legume containing mixes.

Do not roll or harrow either one week pre or post treatment.

Treat only vigorous, healthy crops growing under good soil and weather conditions.

Do not treat diseased crops or those suffering pest attack.

Do not treat crops suffering herbicide damage, drought, waterlogging, other stress.

Cereals undersown with grasses may be treated provided that the grasses are tillering.

### **SPRAY MIXING**

Half fill the spray tank with water. Pour the required volume of FLUXYR 200 EC into the spray tank, then top up with the required amount of water. Stir continuously. Spray volume from 200 to 400l/ha. Use equipment that is clean and in good working order. Avoid overlapping spray swaths. Spray should be applied as a MEDIUM spray (BCPC definition).

## **PRECAUTIONS**

Flash point: 55°C

Freezing point: <-5°C

Keep in original container.

Do not re-use container. The empty container must be sent for disposal as hazardous waste.

Unused product should be disposed of as hazardous waste, in compliance with local and national legislation.

When using the product thoroughly empty and rinse the product container, pouring the rinse water into the spray tank.

Clean sprayer and tank thoroughly after use with a wetting agent.

## **MODE OF ACTION**

Fluroxypyr, the active ingredient of FLUXYR 200 EC, belongs to the family of synthetic auxin herbicides. Repeated use of preparations containing substances with the same mode of action may lead to the emergence of resistant strains. To reduce the risk of occurrence of this phenomenon, it is advisable to avoid the repeated use of products with the same mode of action and alternate applications of products with different modes of action, both on one crop and in rotation of crops.

## **RATE AND TIME OF APPLICATION AND TARGET WEEDS**

For optimum efficacy application of FLUXYR 200 EC to weeds should be when the weeds are actively growing.

| <b>Crop</b>                           | <b>Timing</b>   | <b>Dose</b> | <b>Weed susceptibility</b>  |
|---------------------------------------|---|-------------|---|
| Winter barley,<br>winter wheat        | From the crop having 2 leaves (GS12) until the fully tillered stage (GS29).<br><br>Do not treat crops once they have passed the fully tillered stage (GS29) wait until after the 1 <sup>st</sup> March. | 0.75 l/ha   | Cleavers (to 5cm)<br>Common chickweed (to 5 cm)   |
|                                       | After 1 <sup>st</sup> March from the crop having 2 leaves (GS12) until the flag leaf stage: flag leaf fully unrolled, ligule just visible stage.  | 1.0 l/ha    | Black bindweed (to 6 leaves)<br>Cleavers (to flowering stage)<br>Common chickweed (to flowering stage)<br>Forget-me-not (to flowering stage)<br>Hempnettle (to flowering stage)<br>Knotgrass (to 2 leaves)<br>Red dead-nettle (to 4 leaves) |
| Triticale, winter<br>oats, winter rye | In spring (after 1 <sup>st</sup> March) from the crop having 2 leaves unfolded to before the 2 <sup>nd</sup> node is detectable.  | 1.0 l/ha    | Black bindweed (to 6 leaves)<br>Cleavers (to flowering stage)<br>Common chickweed (to flowering stage)<br>Forget-me-not (to flowering stage)<br>Hempnettle (to flowering stage)<br>Knotgrass (to 2 leaves)<br>Red dead-nettle (to 4 leaves) |
| Spring barley,<br>spring wheat        | From the crop having 2 leaves unfolded to Flag leaf stage: flag leaf fully unrolled, ligule just visible stage.   | 0.75 l/ha   | Black bindweed (to 4 leaves)<br>Cleavers (to 10 cm)<br>Common chickweed (to 10 cm)<br>Forget-me-not (to 5 cm)<br>Hempnettle (to 10 cm)<br>Knotgrass (to 2 leaves)   |
| Spring oats                           | From the crop having 2 leaves unfolded to before the 2 <sup>nd</sup> node is detectable.  | 0.75 l/ha   | Black bindweed (to 4 leaves)<br>Cleavers (to 10 cm)<br>Common chickweed (to 10 cm)<br>Forget-me-not (to 5 cm)<br>Hempnettle (to 10 cm)<br>Knotgrass (to 2 leaves)   |
| Forage maize                          | Between the crop having 3 and 6 leaves unfolded, inclusive. Treat before the buttress roots start to develop.   | 1.0 l/ha    | Black bindweed (to 4 leaves)<br>Cleavers (to 10 cm)<br>Common chickweed (to 10 cm)<br>Forget-me-not (to 5 cm)<br>Hempnettle (to 10 cm)<br>Knotgrass (to 2 leaves)<br>Black nightshade (to 4 leaves)   |

|  |  |   |  |
|--|--|---|--|
| <b>Crop</b><br>Grassland<br>(newly sown leys)    | <b>Timing</b><br>Susceptible annual weeds<br><br>In autumn or spring. From the grasses having at least 3 expanded leaves.  | <b>Dose</b><br>0.75/ha  | <b>Weed susceptibility</b><br>Common Chickweed to 5 cm                                 |
| Important: any clovers will be killed            |  |   |  |
| Established Grassland (permanent and rotational) | Control of Dock spp.<br>Cut the docks before seeding, normally late-May to early-June. Spray the regrowth 2-3 weeks after cutting. A repeat treatment may be required in the following year if the docks are well-established or numerous. When making spot treatments, spray to just wet the dock leaves without run-off. | 2 l/ha (or 30ml FLUXYR 200 EC per 10 litres water spot treatment) | Broad-leaved dock.<br>Curled-leaved dock – both at the rosette stage before flowering. |
| Important: any clovers will be killed            |  |   |  |

#### **CROP FAILURE**

In the event of a cereal or grass crop failure the following crops can be sown: spring cereals, spring oilseed rape, maize, onion, cultivated poppies and new leys. A minimum interval of 5 weeks is required and there are no cultivation restrictions.

#### **ROTATIONAL CROPS**

All crops can be sown following an application of FLUXYR 200 EC.

#### **COMPATIBILITY**

For information on tank mixes consult the UPL website – [ie.uplonline.com](http://ie.uplonline.com)

#### **CONDITIONS OF SUPPLY**

All goods supplied by us are of high grade and we believe them to be suitable but, as we cannot exercise control over their storage, handling, mixing or use, or of the weather conditions before, during and after application, which may affect the performance of the goods, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded, and no responsibility will be accepted by us or resellers for any failure in performance, damage or injury whatsoever arising from their storage, handling, application or use.

These conditions cannot be varied by our staff or agents whether or not they supervise or assist in the use of such goods.

To access the Safety Data Sheet for FLUXYR 200 EC please use the link below or scan the code.



[ie.uplonline.com/cropproductportfolio](http://ie.uplonline.com/cropproductportfolio)

Alternatively, contact your local supplier.

# FLUXYR 200 EC<sup>®</sup>

PCS 05787

An emulsifiable concentrate containing 200g/L fluroxypyr (equivalent to 28.6% w/w fluroxypyr-1-methylheptyl ester with an aromatic hydrocarbon).

For the control of certain annual weeds in the listed cereals, maize and grassland.

FOR PROFESSIONAL USE ONLY

| <b>Crop</b>                        | <b>Maximum individual dose (L/product/ha)</b> | <b>Maximum total dose (L/product/ha)</b> | <b>Latest time of application</b>          |
|------------------------------------|---|--|--|
| Spring wheat, spring barley        | 0.75  | 0.75                                     | Flag leaf ligule just visible (GS39)       |
| Winter wheat, winter barley        | 1.0   | 1.0                                      | Flag leaf ligule just visible (GS39)       |
| Winter rye, winter oats, triticale | 1.0   | 1.0                                      | Before second node detectable stage (GS32) |
| Spring oats                        | 0.75  | 0.75                                     | Before second node detectable stage (GS32) |
| Forage maize                       | 1.0   | 1.0                                      | Before 7 leaves unfolded (GS17)            |
| Grassland (established)            | 2.0   | 2.0                                      | 7 days livestock exclusion                 |
| Grassland (new leys)               | 0.75  | 0.75                                     | 7 days livestock exclusion                 |

**OTHER SPECIFIC RESTRICTIONS**  
A maximum total dose of 0.75 litres per hectare must be observed for applications made to cereals between crop emergence in the year of planting and 1st February in the year of harvest.  
When the product is applied as a spot treatment to permanent grassland and rotational grass, the maximum concentration must not exceed 30ml product per 10 litres water.  
To protect aquatic organisms respect an unsprayed buffer zone of 5m to surface water bodies.

Do not contaminate water with the product or its container. Do not clean application equipment near surface water/avoid contamination via drains from farmyards and roads.

**SHAKE WELL BEFORE USE**  
**PROTECT FROM FROST**

For Batch Number and  
Manufacturing Date see container

**UPL Europe Ltd**  
The Centre, 1st Floor, Birchwood Park, Warrington, WA3 6YN  
**Telephone:** 01925 819999 **Fax:** 01925 817425

**For 24 hour emergency information contact:**  
**CARECHEM24 : +44 (0) 1235 239670**

FLUXYR/IE/SL/B/01117/UPL